

Integration 2

Affine substitution

If $\int f(x) dx = g(x)$ and a, b are constant, then

$$\int f(ax + b) dx = g(ax + b)/a$$

$$\int \cos(x) dx = \sin(x)$$

$$\int \cos(2x + 3) dx = \sin(2x + 3)/2$$

$$\int e^x dx = e^x$$

$$\int e^{-2x+7} dx = e^{-2x+7}/(-2)$$

$$\int \tan(x) dx = -\ln(\cos(x))$$

$$\int \tan(\pi x) dx = -\ln(\cos(\pi x))/\pi$$


Exponential oscillations

- ▶ An *exponential oscillation* is a function of the form

$$f(x) = e^{\lambda x} (a \cos(\omega x) + b \sin(\omega x)),$$

where a , b , λ and ω are constants.

- ▶ The *growth rate* is λ , and the *angular frequency* is ω .



- ▶ Special cases:

$$f(x) = e^{\lambda x} \sin(\omega x) \qquad (a = 0, b = 1)$$

$$f(x) = a \cos(\omega x) + b \sin(\omega x) \qquad (\lambda = 0)$$

$$f(x) = ae^{\lambda x} \qquad (\omega = 0).$$

Integrating exponential oscillations

The integral of an EO is another EO with the same growth rate and angular frequency.

$$\int e^{\lambda x} (a \cos(\omega x) + b \sin(\omega x)) dx = e^{\lambda x} (A \cos(\omega x) + B \sin(\omega x))$$

$$A = \frac{a\lambda - b\omega}{\lambda^2 + \omega^2} \quad B = \frac{a\omega + b\lambda}{\lambda^2 + \omega^2}.$$

► Example: find

$$\int e^{-2x} (5 \cos(4x) - 3 \sin(4x)) dx \quad \int e^{-2x} (5 \cos(4x) - 3 \sin(4x)) dx$$

► $\lambda = -2, \omega = 4, a = 5, b = -3$

► $A = \frac{a\lambda - b\omega}{\lambda^2 + \omega^2} = \frac{5 \cdot (-2) - (-3) \cdot 4}{(-2)^2 + 4^2} = 1/10$

► $B = \frac{a\omega + b\lambda}{\lambda^2 + \omega^2} = \frac{5 \cdot 4 + (-3) \cdot (-2)}{(-2)^2 + 4^2} = 13/10$

$$\int e^{-2x} (5 \cos(4x) - 3 \sin(4x)) dx = e^{-2x} (\cos(4x) + 13 \sin(4x))/10$$

Alternatively:

$$\int e^{-2x}(5 \cos(4x) - 3 \sin(4x)) dx = e^{-2x}(A \cos(4x) + B \sin(4x)) \text{ for some } A, B$$

$$\begin{aligned} e^{-2x}(5 \cos(4x) - 3 \sin(4x)) &= \frac{d}{dx} \left(e^{-2x}(A \cos(4x) + B \sin(4x)) \right) \\ &= -2e^{-2x}(A \cos(4x) + B \sin(4x)) + \\ &\quad e^{-2x}(-4A \sin(4x) + 4B \cos(4x)) \\ &= e^{-2x}((4B - 2A) \cos(4x) - (2B + 4A) \sin(4x)) \end{aligned}$$

By comparing coefficients, we must have $4B - 2A = 5$ and $2B + 4A = 3$. These equations can be solved to give $A = 1/10$ and $B = 13/10$. Thus

$$\int e^{-2x}(5 \cos(4x) - 3 \sin(4x)) dx = e^{-2x}(\cos(4x) + 13 \sin(4x))/10.$$

Polynomial exponential oscillations

- ▶ A *polynomial exponential oscillation* is a function of the form

$$f(x) = e^{\lambda x}(a(x) \cos(\omega x) + b(x) \sin(\omega x)),$$

where $a(x)$ and $b(x)$ are polynomials.

- ▶ λ is the *growth rate* and ω is the *angular frequency*. The *degree* is the highest power of x that occurs in $a(x)$ or in $b(x)$.
- ▶ The function $f(x) = e^{-2x}((1 + x^5) \cos(4x) + x^3 \sin(4x))$ is a PEO of growth rate -2 , frequency 4 and degree 5 .
- ▶ The function $f(x) = e^{4x}((1 + x^3 + x^6) \sin(3x))$ is a PEO of growth rate 4 , frequency 3 and degree 6 .
- ▶ **Fact:** The integral of any PEO is another PEO with the same growth rate, frequency and degree.

Integrating PEO's — I

- ▶ $\int xe^{-x} \sin(x) dx$ is a PEO of degree 1, growth -1 , frequency 1.
- ▶ $\int xe^{-x} \sin(x) dx = (Ax + B)e^{-x} \cos(x) + (Cx + D)e^{-x} \sin(x)$
for some A, B, C, D .
- ▶
$$\begin{aligned}xe^{-x} \sin(x) &= \frac{d}{dx} ((Ax + B)e^{-x} \cos(x) + (Cx + D)e^{-x} \sin(x)) \\ &= Ae^{-x} \cos(x) - (Ax + B)e^{-x} \cos(x) - (Ax + B)e^{-x} \sin(x) + \\ &\quad Ce^{-x} \sin(x) - (Cx + D)e^{-x} \sin(x) + (Cx + D)e^{-x} \cos(x) \\ &= (-A + C)xe^{-x} \cos(x) + (A - B + D)e^{-x} \cos(x) + \\ &\quad (-A - C)xe^{-x} \sin(x) + (-B + C - D)e^{-x} \sin(x).\end{aligned}$$
- ▶ $-A + C = 0, A - B + D = 0, -A - C = 1, -B + C - D = 0$.
- ▶ So $A = -1/2, B = -1/2, C = -1/2, D = 0$
- ▶ $\int xe^{-x} \sin(x) dx = -((x + 1)e^{-x} \cos(x) + xe^{-x} \sin(x))/2$.

Integrating PEO's — II

- ▶ $\int x^3 e^x dx$ is a PEO of degree 3, growth 1 and frequency 0.
- ▶ $\int x^3 e^x dx = (Ax^3 + Bx^2 + Cx + D)e^x$ for some A, B, C, D .
- ▶
$$\begin{aligned}x^3 e^x &= \frac{d}{dx} \left((Ax^3 + Bx^2 + Cx + D)e^x \right) \\ &= (3Ax^2 + 2Bx + C)e^x + (Ax^3 + Bx^2 + Cx + D)e^x \\ &= (Ax^3 + (3A + B)x^2 + (2B + C)x + (C + D))e^x.\end{aligned}$$
- ▶ $A = 1, 3A + B = 0, 2B + C = 0, C + D = 0$.
- ▶ so $A = 1, B = -3, C = 6, D = -6$
- ▶ so $\int x^3 e^x dx = (x^3 - 3x^2 + 6x - 6)e^x$.

Integration by parts — I

▶ Consider $\int xe^{x/a} dx$.

▶ Consider $\int xe^{x/a} dx$.

▶ $u = x$

$$dv/dx = e^{x/a}$$

▶ $du/dx = 1$

$$v = a e^{x/a}$$

▶ $\int xe^{x/a} dx = uv - \int \frac{du}{dx} v dx = axe^{x/a} - \int a e^{x/a} dx = axe^{x/a} - a^2 e^{x/a}$

▶ To integrate a product, call the factors u and $\frac{dv}{dx}$.

▶ Differentiate u to find du/dx .

▶ Integrate $\frac{dv}{dx}$ to find v .

▶ Use the formula:

$$\int u \frac{dv}{dx} dx = uv - \int \frac{du}{dx} v dx$$

▶ This is most useful when (a) du/dx is simpler than u (eg u polynomial) and (b) v is no more complicated than dv/dx (eg $dv/dx = \cos(x)$).

Integration by parts — II

- ▶ Consider $\int (1 - \ln(x))x^{-2} dx$.
 - ▶ Consider $\int (1 - \ln(x))x^{-2} dx$.
 - ▶ $u = 1 - \ln(x)$ $dv/dx = x^{-2}$
 - ▶ $du/dx = -x^{-1}$ $v = -x^{-1}$
 - ▶
$$\int (1 - \ln(x))x^{-2} dx = uv - \int \frac{du}{dx} v dx = -(1 - \ln(x))x^{-1} - \int x^{-2} dx$$
$$= (\ln(x) - 1)x^{-1} + x^{-1} = \ln(x)/x$$
-

- ▶ To integrate a product, call the factors u and $\frac{dv}{dx}$.
- ▶ Differentiate u to find du/dx .
- ▶ Integrate $\frac{dv}{dx}$ to find v .
- ▶ Use the formula:

$$\int u \frac{dv}{dx} dx = uv - \int \frac{du}{dx} v dx$$

Integration by parts — III

- ▶ Consider $\int x \sin(\omega x) dx$.
 - ▶ Consider $\int x \sin(\omega x) dx$.
 - ▶ $u = x$ $dv/dx = \sin(\omega x)$
 - ▶ $du/dx = 1$ $v = -\omega^{-1} \cos(\omega x)$
 - ▶
$$\int x \sin(\omega x) dx = uv - \int \frac{du}{dx} v dx = -\omega^{-1} x \cos(\omega x) + \int \omega^{-1} \cos(\omega x) dx$$
$$= -\omega^{-1} x \cos(\omega x) + \omega^{-2} \sin(\omega x)$$
-

- ▶ To integrate a product, call the factors u and $\frac{dv}{dx}$.
- ▶ Differentiate u to find du/dx .
- ▶ Integrate $\frac{dv}{dx}$ to find v .
- ▶ Use the formula:

$$\int u \frac{dv}{dx} dx = uv - \int \frac{du}{dx} v dx$$

Integration by parts — IV

- ▶ Consider $\int \arcsin(x) dx$.
 - ▶ Consider $\int \arcsin(x) \cdot 1 dx$.
 - ▶ $u = \arcsin(x)$ $dv/dx = 1$
 - ▶ $du/dx = (1 - x^2)^{-1/2}$ $v = x$
 - ▶
$$\int \arcsin(x) \cdot 1 dx = uv - \int \frac{du}{dx} v dx = \arcsin(x) \cdot x - \int x(1 - x^2)^{-1/2} dx$$
$$= x \arcsin(x) + (1 - x^2)^{1/2}$$
-

- ▶ To integrate a product, call the factors u and $\frac{dv}{dx}$.
- ▶ Differentiate u to find du/dx .
- ▶ Integrate $\frac{dv}{dx}$ to find v .
- ▶ Use the formula:

$$\int u \frac{dv}{dx} dx = uv - \int \frac{du}{dx} v dx$$