

De Rham cohomology

Given a suitable space X (for example, an open subset of \mathbb{R}^n) one can define vector spaces $H^k(X)$ for integers $k \geq 0$, called the *de Rham cohomology groups* of X . If X is an open subset of \mathbb{R}^3 , then the definition can be phrased in terms of the usual operations of vector calculus (**div**, **grad** and **curl**), giving a close relationship with some physical phenomena in electromagnetism. The same circle of ideas also gives a nice unified picture of results such as the Fundamental Theorem of Calculus, Green's Theorem, Stokes' Theorem, and the Divergence Theorem. The dimensions of the spaces $H^k(X)$ give information about the topological structure of X , such as the number of holes. With more work, one can generalise everything to open subsets of \mathbb{R}^n for $n > 3$. The main problem is to generalise the definitions of the dot and cross products, a topic known as *exterior algebra*. One also wants to define de Rham cohomology for sets such as curves and surfaces (which are closed, rather than open, in \mathbb{R}^3) and higher-dimensional generalisations (known as *submanifolds* of \mathbb{R}^n).

Another important topic is the multiplicative structure. Given $a \in H^i(X)$ and $b \in H^j(X)$ there is a natural definition of a product $ab \in H^{i+j}(X)$. This makes the vector space $H^*(X) = \bigoplus_i H^i(X)$ into a ring. As well as setting up the general theory, one can give a complete description of $H^*(X)$ for many interesting spaces (such as spheres, the torus, configuration spaces and so on). There are many applications, including the Brouwer Fixed Point Theorem: any continuous map $f: B^n \rightarrow B^n$ has a fixed point (where $B^n = \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\| \leq 1\}$).

REFERENCES

- [1] Ib Madsen and Jørgen Tornehave, *From calculus to cohomology*, Cambridge University Press, Cambridge, 1997. de Rham cohomology and characteristic classes. MR 98g:57040
- [2] Michael Spivak, *Calculus on manifolds. A modern approach to classical theorems of advanced calculus*, W. A. Benjamin, Inc., New York-Amsterdam, 1965. MR 35 #309
- [3] John W. Milnor, *Topology from the differentiable viewpoint*, Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 1997. Based on notes by David W. Weaver; Revised reprint of the 1965 original. MR 98h:57051
- [4] Raoul Bott and Loring W. Tu, *Differential forms in algebraic topology*, Graduate Texts in Mathematics, vol. 82, Springer-Verlag, New York, 1982. MR 83i:57016

Supervisor: Professor Neil Strickland
Room J26, N.P.Strickland@shef.ac.uk

October 14, 2008