

## Soluble groups

A *subnormal series* for a finite group  $G$  is a chain of subgroups

$$\{1\} = H_0 \leq H_1 \leq \cdots \leq H_{r-1} \leq H_r = G$$

such that  $H_{i-1}$  is a normal subgroup of  $H_i$  (for  $i = 1, \dots, r$ ); this means that we can define quotient groups  $H_i/H_{i-1}$ . We say that  $G$  is *soluble* if there exists a subnormal series for which all the quotient groups  $H_i/H_{i-1}$  are abelian.

Soluble groups are easier to handle than more general groups, because many things can be reduced to the abelian case. Solubility is important in Galois theory, because the roots of a polynomial can be expressed in terms of radicals iff the corresponding Galois group is soluble. A very important theorem, far beyond the scope of this project, says that all groups of odd order are soluble; the proof takes hundreds of pages.

This project should start with some generalities about subnormal series. In particular, a *composition series* for a group  $G$  is a subnormal series all of whose quotients are simple groups; these quotients are called the *composition factors* of the group. The Jordan-Hölder theorem says that they are well-defined, despite the fact that  $G$  will typically have many different composition series; this should be proved in the project.

Cyclic and dihedral groups are soluble. The symmetric and alternating groups  $S_n$  and  $A_n$  are soluble for  $n \leq 4$ , but not for  $n \geq 5$ . Subgroups and quotient groups of soluble groups are soluble. A group of order  $p^a$  (with  $p$  prime) is always soluble, by a fairly easy argument. A group of order  $p^a q^b$  (with  $p$  and  $q$  both prime) is also soluble, by a theorem of Burnside, which should be accessible to a student who has studied Representation Theory.

The following theorem of Hall could form the main goal of the project:

**Theorem:** *Suppose that  $G$  is soluble and  $|G| = mn$  with  $m$  and  $n$  coprime; then  $G$  has a subgroup of order  $m$ , and any two such groups are conjugate.*

(The case where  $m$  is a power of a prime is a theorem of Sylow, which holds even when  $G$  is not soluble. When  $m$  is not a power of a prime, solubility is definitely needed.)

## REFERENCES

- [1] John B. Fraleigh, *A first course in abstract algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1967.MR 37 #1212
- [2] Ian D. Macdonald, *The theory of groups*, Clarendon Press, Oxford, 1968.MR 39 #5670
- [3] Jean-Pierre Serre, *Linear representations of finite groups*, Springer-Verlag, New York, 1977. Translated from the second French edition by Leonard L. Scott; Graduate Texts in Mathematics, Vol. 42.MR 56 #8675
- [4] Marshall Hall Jr., *The theory of groups*, Chelsea Publishing Co., New York, 1976. Reprinting of the 1968 edition.MR 54 #2765
- [5] M. Aschbacher, *Finite group theory*, 2nd ed., Cambridge Studies in Advanced Mathematics, vol. 10, Cambridge University Press, Cambridge, 2000.MR 2001c:20001

**Supervisor: Professor Neil Strickland**  
**Room J26, N.P.Strickland@shef.ac.uk**

October 14, 2008